

1. Construct an explicit deformation retraction from  $\mathbb{R}^n \setminus \{0\}$  onto  $S^1$ .
2. Show that a retract of a contractible space is contractible.
3. Show that a space  $X$  is contractible if and only if every map  $f : X \rightarrow Y$  for arbitrary  $Y$  is nullhomotopic.
4. Let  $X$  be a path-connected topological space. Show that  $\pi_1(X)$  is abelian if and only if the base change homomorphism  $\beta_h$  depends only on the endpoint of the path  $h$ .
5. If  $X_0$  is a path component of a topological space  $X$  containing the base point  $x_0$ , show that the inclusion  $X_0 \hookrightarrow X$  induces an isomorphism on  $\pi_1$  (basepoint being  $x_0$ ).
6. Construct infinitely many non-homotopic retractions  $S^1 \vee S^1 \rightarrow S^1$ .
7. Suppose  $f_t : X \rightarrow X$  is a homotopy such that  $f_0$  and  $f_1$  are each the identity map. Show that for any  $x_0 \in X$ ,  $t \mapsto f_t(x_0)$  considered as a loop in  $\pi_1(X, x_0)$  lies in the center.
8. Let  $X$  be the quotient space of  $S^2$  obtained by identifying the north and south poles to a point. Put a cell complex structure on  $X$  and use this to compute its  $\pi_1$ .
9. Show that the join  $X * Y$  of two non-empty spaces  $X$  and  $Y$  is simply connected if  $X$  is path-connected.
10. Let  $\tilde{X}$  and  $\tilde{Y}$  be two simply connected covering spaces of path-connected, locally path-connected spaces  $X$  and  $Y$ . Show that if  $X \cong Y$  then  $\tilde{X} \cong \tilde{Y}$ .
11. Show that if  $A$  is a retract of  $X$  then the map  $H_n(A) \rightarrow H_n(X)$  induced by the inclusion is injective.
12. Show that  $H_0(X, A) = 0$  iff  $A$  meets every path component of  $X$ .
13. Show that  $H_1(X, A) = 0$  if and only if  $H_1(A) \rightarrow H_1(X)$  is surjective and each path component of  $X$  contains at most one path component of  $A$ .
14. Show that for the subspace  $\mathbb{Q} \in \mathbb{R}$  the relative homology group  $H_1(\mathbb{R}, \mathbb{Q})$  is a free abelian group. Find a basis.
15. Describe a CW structure on the following spaces and compute their homology:
  - (a)  $\mathbb{C}P^n$ .
  - (b)  $M_g$ : a closed orientable surface of genus  $g$ .
  - (c)  $\mathbb{R}P^n$ .
16. If a finite CW complex  $X$  is a union of subcomplexes  $A$  and  $B$  then show that  $\chi(X) = \chi(A) + \chi(B) - \chi(A \cap B)$ .
17. Given a map  $f : S^{2n} \rightarrow S^{2n}$  show that there exists some point  $x \in S^{2n}$  with either  $f(x) = x$  or  $f(x) = -x$ . Deduce that every map  $\mathbb{R}P^{2n} \rightarrow \mathbb{R}P^{2n}$  has a fixed point.
18. Let  $M_g$  denote the surface of genus  $g$  with its standard embedding in  $\mathbb{R}^3$ . Let  $R$  denote the compact region in  $\mathbb{R}^3$  that it bounds. Let  $X$  denote two copies of  $R$  glued along the boundary surface  $M_g$ . Compute the homology groups of  $X$  via the Mayer-Vietoris sequence for this decomposition of  $X$  into two copies of  $R$ .